CONTENTS

	Page
Meat Extracts, Their Composition and Identification. JAMES A. EMERY and ROBERT R. HENLEY	1
Quantity and Composition of Ewes' Milk: Its Relation to the Growth of Lambs. RAY E. NEIDIG and E. J. IDDINGS	19
Seed Disinfection by Formaldehyde Vapor. CECIL C. THOMAS	33
Influence of Soil Environment on the Rootrot of Tobacco. JAMES JOHNSON and R. E. HARTMAN	41
Relation of Sulphates to Plant Growth and Composition. H. G. MILLER	87
Relation of Weather to Fruitfulness in the Plum. M. J. DORSEY	103
Structure of the Maize Ear as Indicated in Zea-Euchlaena Hybrids. G. N. COLLINS	127
Carbohydrate Metabolism in Green Sweet Corn during Storage at Different Temperatures. CHARLES O. APPLEMAN and JOHN M. ARTHUR	137
Certain Relationships between the Flowers and Fruits of the Lemon. HOWARD S. REED	153
Ultra-Microscopic Examination of Disperse Colloids Present in Bituminous Road Materials. E. C. E. LORD	167
Derris as an Insecticide. N. E. MCINDOO, A. F. SEIVERS, and W. S. ABBOTT	177
Effects of Heat on Trichinæ. B. H. RANSOM and BENJAMIN SCHWARTZ	201
Effect of Removing the Pulp from Camphor Seed on Germination and the Subsequent Growth of the Seedlings. G. A. RUSSELL	223
Bacterium abortus Infection of Bulls. J. M. BUCK, G. T. CREECH, and H. H. LADSON	239
Investigations on the Mosaic Disease of the Irish Potato. E. S. SCHULTZ, DONALD FOLSOM, F. MERRILL HILDEBRANDT, and LON A. HAWKINS	247
Temperature in Relation to Quality of Sweetcorn. NEIL E. STEVENS and C. H. HIGGINS	275
Variation of Ayrshire Cows in the Quantity and Fat Content of Their Milk. RAYMOND PEARL and JOHN RICE MINER	285
Index	323

ERRATA AND AUTHORS' EMENDATIONS

Page 82, Table II, "Gm." should read "Pounds."

Page 158, line 2, "variability" should read "viability."

Page 160, line 23, "0.183" should read "−0.183."

Page 162, Table VII, "Months required for maturity" should stand above the table. The legend at the left of the table should be "Month in which fruit set."

Page 179, line 30, "petroleum, ether" should read "petroleum ether."

ILLUSTRATIONS

PLATES

Influence of Soil Environment on the Rootrot of Tobacco

Page

Plate 1. I.—Influence of amount of infestation on injury from tobacco rootrot: A, All uninfested soil; B, three-fourths uninfested soil; C, one-half uninfested soil; D, one-fourth uninfested soil; E, all infested soil. II, III.—Influence of moisture content of soil on the amount of injury done by the tobacco rootrot; II, infested soil; III, uninfested soil (control series)—1A, one-fourth saturation infested soil; 2A, one-half saturation infested soil; 3A, three-fourths saturation infested soil; 4A, full saturation infested soil; 1B, one-fourth saturation uninfested soil; 2B, one-half saturation uninfested soil; 3B, three-fourths saturation uninfested soil; 4B, full saturation uninfested soil. IV.—Influence of soil temperature on the growth of tobacco in infested soil (jars to left of temperature labels) and in uninfested soil (jars to right of temperature labels) at temperatures of approximately 13°, 17°, 23°, 26°, and 36° C. 86

Plate 2. I.—Soil temperature tanks used in the temperature experiments. II, III.—Influence of soil temperature on the growth of tobacco: 1A, infested soil, 17°–18° C.; 2A, infested soil, 20°–21° C; 3A, infested soil, 23°–24° C; 4A, infested soil, 25°–26° C.; 5A, infested soil, 28°–29° C.; 6A, infested soil, 31°–32° C.; 1B, uninfested soil, 17°–18° C.; 2B, uninfested soil, 20°–21° C.; 3B, uninfested soil, 23°–24° C.; 4B, uninfested soil, 25°–26° C.; 5B, uninfested soil, 28°–29° C.; 6B, uninfested soil, 31°–23° C. IV.—Influence of different soil temperatures on root development: 1A, uninfested soil, 17°–18° C.; 1B, infested soil, 17°–18° C.; 2A, uninfested soil, 20°–21° C.; 2B, infested soil, 20°–21° C.; 3A, uninfested soil, 23°–24° C.; 3B, infested soil, 23°–24° C.; 4A, uninfested soil, 25°–26° C.; 4B, infested soil, 25°–26° C.; 5A, uninfested soil, 28°–29° C.; 5B, infested soil, 28°–29° C; 6A, uninfested soil, 31°–32° C.; 6B, infested soil, 31°–32° C. 86

Plate 3. Influence of high (30° C.) and low (20° C.) soil-temperature on recovery of plants in infested soil. 86

Plate 4. I, II.—Influence of soil reaction on extent of damage by tobacco rootrot: I, Infested soil; II, uninfested soil—1A, infested soil, lime requirement 9.38 tons per acre; 2A, infested soil, lime requirement 7.19 tons per acre; 3A, infested soil, lime requirement 4.60 tons per acre; 4A, infested soil, lime requirement 2.62 tons per acre; 5A, infested soil, lime requirement 0.72 ton per acre; 6A, infested soil, slightly alkaline; 7A, infested soil, strougly alkaline; 1B, uninfested soil, lime requirement 9.38 tons per acre; 2B, uninfested soil, lime requirement 7.19 tons per acre; 3B, uninfested soil, lime requirement 4.60 tons per acre; 4B, uninfested soil, lime requirement 2.62 tons per acre; 5B, uninfested soil, lime requirement 0.72 ton per acre; 6B, uninfested soil, slightly alkaline; 7B, uninfested soil, strongly alkaline. III, IV.—Influence of the amount of organic matter in the soil on injury by tobacco rootrot: III, 1A–6A, Influence of gradually increasing amounts of organic matter in infested soil from 1A, no organic matter, to 6A, all leaf mold. IV.—1B–6B, Influence of gradually increasing amounts of organic matter in uninfested siol from 1B, no organic matter, to

Page

6B, all leaf mold (control series). V, VI.—Influence of the amount of organic matter in the soil on injury by tobacco rootrot: V, 1A–5A, Influence of gradually increasing amounts of organic matter in uninfested soil from 1A, no organic matter, to 5A, all leaf mold (control series); VI, 1B–5B, Influence of gradually increasing amounts of organic matter in infested soil from 1B, no organic matter, to 5B, all leaf mold..... 86

PLATE 5. I.—Influence of relative amount of sand and clay on tobacco rootrot: A, uninfested series: B, infested series—1A, uninfested soil, three-fourths clay and one-fourth sand; 1B, infested soil, three-fourths clay and one-fourth sand; 2A, uninfested soil, one-half clay and one-half sand; 2B, infested soil, one-half clay and one-half sand; 3A, uninfested soil, three-fourths sand; 3B, infested soil, three-fourths sand; 4A, uninfested soil, all sand; 4B, infested soil, all sand. II, III.—Influence of soil fertility on amount of tobacco rootrot: II, infested series; III, uninfested series—1A, infested soil, no treatment; 2A, infested soil, 3.5 gms. of nutrient salts; 3A, infested soil, 7.0 gms. of nutrient salts; 4A, infested soil, 14.00 gms. of nutrient salts; 5A, infested soil, 28 gms. of nutrient salts; 6A, infested soil, 56 gms. of nutrient salts. Note increasing injury from nutrient salts beginning at pot 3A. 1B, uninfested soil, no treatment; 2B, uninfested soil, 3.5 gms. of nutrient salts; 3B, uninfested soil, 7.0 gms. of nutrient salts; 4B, uninfested soil, 14.00 gms. of nutrient salts; 6B, uninfested soil, 28 gms. of nutrient salts; 6B, uninfested soil, 56 gms. of nutrient salts. Note injury from nutrient in pots 5B and 6B. IV.—Relation of compactness of soil to injury caused by *Thielavia basicola:* 1A, infested soil, loosely packed; 1B, uninfested soil, loosely packed; 2A, infested soil, very compact; 2B, uninfested soil, very compact. V.—Influence of transplanting infected seedlings in healthy soil: A, Pennsylvania Broadleaf infected seedlings; B, Pennsylvania Broadleaf healthy seedlings C, White Burley infected seedlings; D. White Burley healthy seedlings; E, Northern Hybrid (a resistant type) infected seedlings; F, Northern Hybrid (a resistant type) healthy seedlings..... 86

PLATE 6. Soil temperature graphs for the month of June, 1915–1918, inclusive, at depths of 2, 4, and 8 inches..... 86

PLATE 7. Soil temperature graphs for the month of July, 1915–1918, inclusive, at depths of 2, 4, and 8 inches..... 86

PLATE 8. Soil temperature graphs for the month of August, 1915–1918, inclusive, at depths of 2, 4, and 8 inches..... 86

RELATION OF SULPHATES TO PLANT GROWTH AND COMPOSITION

PLATE 9. A.—Clover on soil A. The top row, reading from left to right, shows the soil pots which received the following fertilizers: Pot 1, calcium sulphate, sodium nitrate; pot 2, sodium sulphate, sodium nitrate; pot 3, sulphur, sodium nitrate, calcium carbonate; pot 4, sodium nitrate; pot 5, sodium nitrate, calcium carbonate; pot 6, no fertilizer; pot 7, calcium sulphate, sodium nitrate; pot 8, sodium sulphate, sodium nitrate; pot 9, sulphur, calcium carbonate, sodium nitrate; pot 10, sodium nitrate. B.—Clover on soil B. C.—Clover on soil C..... 102

PLATE 10. A.—Rape on soil A. B.—Rape on soil B. C. Rape on soil C.... 102

PLATE 11. A.—Oats on soil A. B.—Oats on soil B. C.—Oats on soil C..... 102

PLATE 12. A.—Oats on sand cultures from soil A. B.—Oats on sand cultures from soil B. C.—Oats on sand cultures from soil C..... 102

RELATION OF WEATHER TO FRUITFULNESS IN THE PLUM

Page

PLATE 13. Plum tree and fruiting branch showing difference between number of flowers borne and quantity of fruit set: A.—The appearance of a plum tree bearing a normal crop of bloom. B.—A single fruiting branch 2 years old showing the contrast to A..... 126

PLATE 14. A.—Stigma of Minnesota No. 21, a greenhouse tree, 24 hours after being selfed, showing the condition of papillate cells in the stigma, pollen tubes, and also traces of the stigmatic fluid. B.—Stigma of Minnesota No. 35, open to cross pollination, showing the condition of a stigma three days after bloom, having withstood a rain of 0.87 inch which fell in the two days previous, lasting in all 18 hours. C.—The turgid papillate cells in Sapa before receptiveness. D.—Opata. Same as C. E.—Abscission layer Minnesota No. 35, showing the cells of the layer 11 days after bloom. F.—The surface at the abscission layer of Assiniboin after the style has fallen, 12 days after bloom..... 126

PLATE 15. Graphic analysis of the weather from the standpoint of wind, sunshine, rain, and temperature for seven years from 1912 to 1918..... 126

STRUCTURE OF THE MAIZE EAR AS INDICATED IN ZEA-EUCHLAENA HYBRIDS

PLATE 16. Intermediate stages between a simple spike of the pistillate inflorescence of Euchlaena and an ear of maize: A.—Spike of pure Florida teosinte. B.—Spike with slightly shortened axis. C.—A still more compact spike with an increased number of seeds. A–C have single spikelets and separate two-ranked alicoles. D.—Spike with single spikelets and yoked alicoles, irregularly four-rowed. E.—Compact spike with two-ranked separate alicoles and single spikelets. F.—Spike with paired spikelets and four ranks of yoked alicoles. G.—Transition stage between four-rowed and eight-rowed ear. H.—Ear of maize with eight rather poorly defined rows of seeds..... 136

PLATE 17. Pistillate inflorescences of hybrid between Euchlaena and maize: A.—Showing pedicelled staminate spikelets with sessile pistillate spikelets. B.—Closely compacted inflorescense with two rows of alicoles and four rows of seeds. C–E.—Spirally twisted inflorescences, with three rows of alicoles..... 136

PLATE 18. Pistillate inflorescences of hybrid between Euchlaena and maize, showing yoked alicoles: A–C.—The alicoles are in four rows corresponding to an eight-rowed ear. D.—The alicoles are in five rows, corresponding to a ten-rowed ear..... 136

ULTRA-MICROSCOPIC EXAMINATION OF DISPERSE COLLOIDS PRESENT IN BITUMINOUS ROAD MATERIALS

PLATE 19. A.—Microscope with ray filter and arc lamp for dark field illumination. B.—Photomicrograph of cross-line micrometer scale, showing colloids in dark field. x320. Taken by E. A. Shuster, jr., Photographic Laboratory, United States Geological Survey..... 176

EFFECT OF REMOVING THE PULP FROM CAMPHOR SEED ON GERMINATION AND SUBSEQUENT GROWTH OF THE SEEDLINGS

PLATE 20. A camphor seed bed, showing the growth of seedlings from pulped and unpulped camphor seed planted in alternate rows..... 238

PLATE 21. A.—Camphor seedlings at the time of transplanting. B.—Camphor seedlings cut back and trimmed ready for transplanting..... 328

BACTERIUM ABORTUS INFECTION OF BULLS

Page

PLATE 22. Photograph of normal and diseased seminal vesicles of bull 98, showing the marked increase in size and the gross pathological changes of one of the organs 246

PLATE 23. A.—Photomicrograph of a section from a normal seminal vesicle of bull. B.—Photomicrograph of section from seminal vesicle of bull 409, showing inflammatory changes 246

PLATE 24. A.—Photomicrograph of section from seminal vesicle of bull 98, showing tissue proliferation and exfoliation of epithelium lining acini. B.—Photomicrograph of section from seminal vesicle of bull 98, showing advanced pathological changes with cell degeneration and necrosis 246

INVESTIGATIONS ON THE MOSAIC DISEASE OF THE IRISH POTATO

PLATE A. Foliage of Irish potato, Green Mountain variety 274

PLATE B. Foliage of potato, Bliss Triumph variety 274

PLATE 25. Leaf of Irish potato, Green Mountain variety, infected with mosaic. Medium stage of disease. 274

PLATE 26. A.—Healthy scion grafted upon diseased stock. B.—An illustration of a method used for introducing aphids 274

PLATE 27. A.—Leaves from graft shown in B of this plate: At right, from healthy parent of scion; at left, from mosaic stock; in center, from mosaic scion. B.—At left, healthy scion grafted to diseased stock, Green Mountain variety; at right, two mosaic shoots of stock. C.—Leaves from corresponding parts of plants shown in Plate 29, B 274

PLATE 28. A.—491x, inoculated artificially with unfiltered juices from mosiac plant February 22 to March 22, 1919. B.—473y, inoculated in same way as 491x, but with juices from healthy plant. 458y, also inoculated with juices from healthy plant 274

PLATE 29. A.—Mosaic of potato transmitted by aphids. 142a, infected plant, Green Mountain variety. B.—Two plants from the same tuber treated alike, except that about 200 aphids were introduced upon one when it was 2 inches high 274

PLATE 30. A.—Inoculated by means of artificial transfers of aphids from diseased plants. Green Mountain variety. B.—Plants inoculated in same way as those in A of this plate, but with aphids taken from healthy plants .. 274

TEXT FIGURES

Page

SEED DISINFECTION BY FORMALDEHYDE VAPOR

FIG. 1. Formaldehyde-vapor disinfecting apparatus..... 34

INFLUENCE OF SOIL ENVIRONMENT ON THE ROT OF TOBACCO

FIG. 1. Soil-thermograph records showing the influence on soil temperature of the shading of soil by growing tobacco..... 68

2. Soil-thermograph records given comparison of a typical record of regulated soil temperature in tanks with a typical record from the field at a depth of 4 inches..... 69

RELATION OF WEATHER TO FRUITFULNESS IN THE PLUM

FIG. 1. An outline drawing of an anther of Minnesota No. 12, showing the adjustment which takes place as a result of taking up or giving off water: A, an anther which has been open in the orchard for three days; B, the same with the anthers pushed up to show the dead area at the upper end of the filament; C, the appearance of the anther after two minutes in water..... 111

STRUCTURE OF THE MAIZE EAR AS INDICATED IN ZEA-EUCHLAENA HYBRIDS

FIG. 1. Diagram showing arrangement of pedicelled and sessile spikelets in A, undifferentiated four-rowed branch; B, eight-rowed ear, the result of the fasciation of two undifferentiated branches; C, eight-rowed ear the result of twisting a single undifferentiated branch; D, 16-rowed ear, the result of fasciation; E, 16-rowed ear, the result of a further twisting of "C"..... 129

CARBOHYDRATE METABOLISM IN GREEN SWEET CORN DURING STORAGE AT DIFFERENT TEMPERATURES

FIG. 1. Depletion of total sugars in green sweet corn during consecutive 24-hour periods of storage at different temperatures..... 146

2. Depletion of sucrose in green sweet corn during consecutive 24-hour periods of storage, expressed as percentages of the initial sucrose in the corn, which was 3.87 per cent, wet weight..... 147

CERTAIN RELATIONSHIPS BETWEEN THE FLOWERS AND FRUITS OF THE LEMON

FIG. 1. Average monthly production of lemon buds during the year..... 155

ULTRA-MICROSCOPIC EXAMINATION OF DISPERSE COLLOIDS PRESENT IN BITUMINOUS ROAD MATERIALS

FIG. 1. Glass slide with ultra-microscope cell drawn to natural scale..... 170

EFFECT OF REMOVING THE PULP FROM CAMPHOR SEED ON GERMINATION AND THE SUBSEQUENT GROWTH OF THE SEEDINGS

Page

FIG. 1. Diagram showing percentage of germination of camphor seed secured from parent tree A under varying conditions 227
2. Graphs showing time required for pulped and unpulped camphor seed to reach maximum germination 229
3. Graph showing time required for camphor seed secured from parent tree A at various times and under various conditions to reach maximum germination 230
4. Diagram showing percentage of total germination of pulped and unpulped camphor seed from 10 parent trees 233
5. Graphs showing rates and percentage of germination of pulped and unpulped camphor seed from 10 parent trees 234

TEMPERATURE IN RELATION TO QUALITY OF SWEETCORN

FIG. 1. Mean hourly temperature for August at Baltimore, Md., and for September, 1918, at Portland, Me 282

VARIATION OF AYRSHIRE COWS IN THE QUANTITY AND FAT CONTENT OF THEIR MILK

FIG. 1. Histograms and fitted curves for variation in mean weekly milk yield of Ayrshire cows of ages 3 to 7 years 306
2. Histograms and fitted curves for variation in mean weekly milk yield of Ayrshire cows of ages 8 to 12 years 307
3. Histograms and fitted curves for variation in fat percentage of milk of Ayrshire cows of ages 3 to 7 years 308
4. Histograms and fitted curves for variation in fat percentage of milk of Ayrshire cows of ages 8 to 12 years 309
5. Showing the change in mean weekly yield of milk in Ayrshire cows. The smooth curve is of the form $yKa=bx=cx^2=d \log x$ 316
6. Showing the observed (zigzag line) and calculated (straight line) changes in the mean fat percentage of the milk of Ayrshire cows with advancing age 318